Saskatchewan
Ministry of
Agriculture

Crop Report

Crop Report for the Period July 27 to August 2, 2010

Published by the Ministry of Agriculture
ISSN 0701 7085
Report number 18, August 2, 2010

Saskatchewan producers have 82 per cent of the hay crop cut and 60 per cent baled or put into silage, according to Saskatchewan Agriculture's weekly Crop Report.

The quality of the hay crop is rated as 17 per cent excellent, 63 per cent good and 10 per cent fair. Haying has progressed furthest in the south, where 86 per cent of the crop has been cut and 68 per cent baled or put into silage. The central regions of the province have 75 per cent of the crop cut and 50 per cent baled or put into silage, and the northern regions have 79 per cent cut and 55 per cent baled or put into silage. High humidity and rain showers continue to slow haying progress. Thunderstorms continue in parts of the province, dropping hail and large amounts of rain.

Harvest has just begun in the southern regions. Fall rye is being swathed and winter wheat combined. Crops are one to two weeks behind normal in development in most areas of the province.

Cropland topsoil moisture is rated as 22 per cent surplus, 72 per cent adequate and six per cent short. Topsoil moisture on hay land and pasture is rated as 14 per cent surplus, 79 per cent adequate, seven per cent short.

The majority of crop damage is being caused by flooding and hail. Insects, disease, gophers and wind are also causing crop damage.

Farmers are busy haying, scouting fields and controlling crop diseases and insects.

One year ago

Eighty-seven per cent of the hay crop was cut, with 68 per cent of this baled or put into silage. Sixteen per cent of the fall rye had been swathed. Combining had started on winter cereals, peas and lentils.

For further information, contact: Grant McLean, PAg,
Cropping Management Specialist, Agriculture Knowledge Centre,
Toll Free: 1-866-457-2377, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.agriculture.gov.sk.ca.

Agriculture and Agri-Food Canada | Agriculture et Agroalimentaire Canada

SCIC
SASKATCHEWAN CROP INSURANCE CORPORATION

Southeastern Saskatchewan (Crop Districts 1, 2, and 3ASE)

Almost all areas in the region reported rain during the past week. The Manor area received 18 mm of rain, the Lampman area 2 mm, the Grenfell area 72 mm, the Wawota area 1 mm, the Griffin area 23 mm, the Odessa area 31 mm, the Indian Head area 11 mm and the Ceylon area 38 mm. The Windthorst and Wilcox areas recorded no rain for the week. Significant rainfall in the Summerberry area filled creeks again. Strong winds and hail were reported in some areas.

Eighty-four per cent of the hay has been cut, with 61 per cent of that baled or put into silage. Sixteen per cent is still standing. Quality is rated as 28 per cent excellent, 52 per cent good and eight per cent fair. Rain throughout the area has slowed haying for some producers. The swaths have been slow to dry due to high humidity and frequent rain showers.

Harvest has just nicely begun. Five per cent of the fall rye has been swathed. A few fields of winter wheat, canola and peas are being cut as well.

Topsoil moisture conditions on cropland are reported as 16 per cent surplus, 80 per cent adequate and four per cent short. On hay land and pasture, topsoil moisture is rated as 13 per cent surplus, 83 per cent adequate and four per cent short. Crop District1B is reporting 25 per cent surplus topsoil moisture on cropland.

Crop damage is mostly attributed to hail and wind. The following communities reported hail damage: Carlyle, Frobisher, Alameda, Manor, Stoughton, Indian Head and Ceylon. Hail stones that fell at Carlyle were jagged and two to four inches in diameter. Leaf diseases are also causing crop damage. Wheat midge were reported in areas of CDs 1B and 2B.

Crops have advanced nicely with the warm weather the past few weeks. Haying is proving difficult, with thunderstorms and high humidity slowing work. Hay quality has been reduced in some areas due to rain on the swaths. Crops are in various stages of development from flowering to ripening. Many areas are still one to two weeks behind normal in crop development. Producers in other areas will start to swath canola and desiccate peas by the end of the week. Weed control on unseeded acres and disease control continues when the weather allows. Farmers are busy haying, controlling weeds in chem-fallow, controlling disease, getting ready for harvest and hauling grain.

Southwestern Saskatchewan (Crop Districts 3ASW, 3AN, 3B and 4)

Thunderstorms and rain were recorded for most of the region, with the southern parts receiving the most rain. The Glentworth area received 60 mm of rain, the Rockglen area 10 mm, the Viceroy area 3 mm, the Mossbank and Coderre areas

4 mm, the Shaunavon area 40 mm, the Mankota area 11 mm, the Kyle area 50 mm, the Gull Lake are 24 mm and the Leader area 19 mm. The Rush Lake, Swift Current and Cabri areas reported no rain for the week.
The region has 89 per cent of the hay cut, with 76 per cent of that baled or put into silage. Eleven per cent is still standing. Quality is rated as 28 per cent excellent and 63 per cent good.

Harvest has just begun, with 15 per cent of the fall rye swathed. Crop reporters are indicating there are a few pea fields ready to straight-combine.

Cropland topsoil moisture is rated as one per cent surplus, 82 per cent adequate, 16 per cent short and one per cent very short. Hay land and pasture topsoil moisture conditions are 84 per cent adequate, 12 per cent short and four per cent very short. Cropland topsoil moisture conditions in CDs 3ASW and 3BS are 33 and 21 per cent short, respectively..

Crop damage has been caused by hail, grasshoppers and gophers. Hail was reported in the following areas: Bengough, Glentworth, Limerick, Lafleche, Hazenmore, Admiral, Consul, Gull Lake and Maple Creek. Spotty and intense rain and hail storms are causing hail damage from 30 to 100 per cent. Crops in some areas could use some rain to help with filling. A few of those areas received the rain this past week, while others continue to miss it. Gophers are causing the most damage in CD 3BS. Grasshoppers are being controlled in lentils.

Farmers are busy haying, scouting fields and controlling weeds on fields that were not seeded. Crop staging varies from field to field. Wheat is heading, canola and mustard are podding and peas and lentils are turning. Harvest should start by the end of the week on early-seeded peas and canola. Later-seeded crops are still flowering. Livestock producers are not cutting too far ahead of the baler to avoid having their hay rained upon.

East-Central Saskatchewan (Crop Districts 5 and 6a)

Thunderstorms throughout the region brought more rain. Most areas reported less than 15 mm of rain, but amounts ranged from nil to 34 mm. The Stockholm, Ituna and Bethune areas received 10 mm, the Saltcoats area 14 mm, the Kuroki area 28 mm, the Craik area 34 mm and the Meacham area 30 mm.

Progress was made with haying during the past week as many areas received only small amounts of rain. Seventy-five per cent of the hay crop is cut and 51 per cent of that has been baled or put into silage; 25 per cent still standing. Frequent rain, wet hay fields and high humidity are slowing progress in some areas. Hay quality is reported as 13 per cent excellent, 66 per cent good and 13 per cent fair. Haying progress is variable across the region. Crop District 5A is reporting 76 per cent is cut (53 per cent baled), CD 5B is reporting 61 per cent

cut (37 per cent baled) and CD 6A is reporting 84 per cent cut (59 per cent baled).

A few fields of winter wheat and fall rye have been swathed.

Topsoil moisture on cropland is reported as 39 per cent surplus, 60 per cent adequate and one per cent short. Hay land and pasture topsoil moisture is rated as 28 per cent surplus, 71 per cent adequate and one per cent short. CD 5B is reporting 76 and 54 per cent surplus topsoil moisture on cropland and hay land and pasture, respectively.

Flooding continues to cause the majority of crop damage. Hail damage was reported in the Saltcoats, Jedburgh, Kamsack and Foam Lake areas.

Crop staging and condition varies across the region. Crop stages ranges from flowering to beginning to turn. There were a couple reports of farmers cutting the weeds in the unseeded acres to try to get the land into shape for next year. Spraying and tilling continues on unseeded acres. The warm weather has pushed crops along.

West-Central Saskatchewan (Crop Districts 6B and 7)

The region recorded rain in amounts ranging from nil to 51 mm. The Dinsmore area received 30 mm of rain, the Langham and Netherhill areas 2 mm, the Biggar area 47 mm, the Unity area 51 mm, the Luseland area 33 mm, the Major, Macklin and Hanley areas received 10 mm.

Seventy-six per cent of the hay crop has been cut and 49 per cent of that has been baled up or put into silage; 24 per cent is still standing. Quality is rated 11 per cent excellent, 59 per cent good and 22 per cent fair. The hay yields look good. The amount of rain on the swath will determine the quality. Hay swaths are slow to dry due to frequent rain and high humidity.

Cropland topsoil moisture is rated as 20 per cent surplus, 75 per cent adequate and five per cent short. Hay land and pasture topsoil moisture conditions are rated as 15 per cent surplus, 77 per cent adequate and eight per cent short.

Most fall rye crops are ready to swath.

The majority of crop damage is due to flooding, wheat midge and grasshoppers. Wheat midge numbers are quite high in areas, and farmers are applying insecticides for control. Grasshopper infestations are not severe in the majority of crops, but farmers are finding sufficient severity in lentil crops to warrant control. Leaf diseases are also causing damage. Hail damage was reported in the Kerrobert area. The heavy rain in some areas has caused some crops to lodge. Crop damage from late-season disease infestations is being reported.

Farmers are busy haying, controlling weeds, disease and insects and hauling grain. Crops are looking pretty good in most areas, although one to two weeks behind normal in development. Crop staging is variable. Producers are looking for some heat to push crops along, especially in lentils where the relatively cool weather has not initiated the end of flowering. Most crops are flowering, while others are podding and starting to fill. For the most part, the crops look good.

Northeastern Saskatchewan (Crop Districts 8 and 9AE)

All areas reported rain from thunderstorms this past week. The Tisdale area received 12 mm of rain, the Arborfield, Codette and Garrick areas recorded 3 mm, the Humboldt area 30 mm, the Lake Lenore and Bruno areas 6 mm and the Melfort area 10 mm.

Eighty-one per cent of the hay crop has been cut, and 57 per cent of that has been baled or put into silage; 19 per cent of the hay crop is still standing. Quality is rated as 85 per cent good. Crop reporters are indicated some hay fields are saturated and some swaths are in the water.

Cropland topsoil moisture is rated as 52 per cent surplus, 46 per cent adequate and two per cent short. Hay land and pasture topsoil moisture is rated as 42 per cent surplus, 54 per cent adequate and four per cent short.

Crop damage is due to flooding and disease. Hail damage was reported in the Tisdale and Nipawin areas. Crops are lodging in areas that received heavy rain.

Farmers are busy haying, controlling weeds and diseases and scouting for insect damage. Wheat midge infestations were heavy in some areas. Crop staging is variable across the region and crops are still one to two weeks behind normal in development. Canola is finishing up flowering and cereals are in the heading stage. Field work continues to be a challenge with the wet fields and continued rain.

Northwestern Saskatchewan (Crop Districts 9AW and 9B)

Thunderstorms rolled through the region last week, leaving rain in amounts ranging from nil to 32 mm. The Leask area received 32 mm, the Spiritwood and Rapid View areas 13 mm, the St. Walburg, Frenchman Butte and Meadow Lake areas received 4 mm and the Dorintosh area 5 mm. The Radisson, Medstead and Glaslyn areas recorded no rain for the week.

Topsoil moisture conditions for cropland are six per cent surplus, 93 per cent adequate and one per cent short. Pasture and hay land conditions are 100 per cent adequate.

Seventy-eight per cent cut of the hay crop has been cut and 54 per cent of that baled or put into silage; 22 per cent is still standing. Quality is reported as six per cent excellent, 69 per cent good and 19 per cent fair. The recent rains have lowered the quality of hay in some areas. Hay yields and pastures look good.

Cropland topsoil moisture is rated as one per cent surplus, 93 per cent adequate and six per cent short. Hay land and pasture topsoil moisture is rated as 93 per cent adequate and seven per cent short.

Wind was the major cause of crop damage.

The warmer temperatures helped advance crops. Most canola and pea crops have finished flowering and are in the podding stage. Cereals have headed out and started to fill. Most crops are one to two weeks behind normal in development, but look quite good. Farmers are busy haying, scouting fields and controlling insects.

Crop Districts and Rural Municipalities in Saskatchewan



Weekly Rainfall

for the week ending August 2, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas this map should be used for regional analysis only

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13, Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 4, 2010

Weekly Rainfall

for the week ending August 2, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 4, 2010

Weekly Rainfall Summary

(in millimeters) 1 inch = 25 mm for the period July 27 to August 2, 2010

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	1	Argyle	N/A	355.0
	2	Mount Pleasant	4	285
	3	Enniskillen	9	292
	33 A	Moose Creek	N/A	249
	33 B	Moose Creek	5	321
	34	Browning	2	254
	61	Antler	N/A	368
	63	Moose Mountain	19	323
	64	Brock	5	328
	65	Tecumseh	8	349
1B	91	Maryfield	26	334
	92	Walpole	1	299
	93	Wawken	11	376
	95	Golden West	NIL	303
	122	Martin	N/A	385
	123 A	Silverwood	6	275
	123 B	Silverwood	N/A	N/A
	124	Kingsley	24	276
	125	Chester	N/A	279
	151 A	Rocanville	9	368
	151 B	Rocanville	22	409
	153	Willowdale	8	407
	154	Esapo	N/A	369
	155	Wolseley	72	462
2A	66	Griffin	23	380
	67	Weyburn	N/A	350
	68	Brokenshell	7	293
	69	Norton	N/A	313
	96	Fillmore	14	264
	97	Wellington	8	219
2B	127 A	Francis	39.5	316
	127 B	Francis	13	213
	129	Bratt's Lake	NIL	233
	156 A	Indian Head	11	389
	156 B	Indian Head	27	367
	157	South Qu'appelle	N/A	NIL
	159	Pense	N/A	304
	161	Moose Jaw	N/A	295
	191	Marquis	N/A	362
3ASE	38 A	Laurier	N/A	337
	38 B	Laurier	6	290
	39 A	The Gap	9	263
	39 B	The Gap	38	407
3ASW	10	Happy Valley	N/A	316
	11	Hart Butte	N/A	NIL
	12	Poplar Valley	9	238
	40	Bengough	47	320
	42	Willow Bunch	13	311
	43	Old Post	10	349
	44	Waverley	60	312
	70	Key West	N/A	239
	71 A	Excel	3	291
	71 B	Excel	4	227
	73 A	Stonehenge	13.5	281
	73 B	Stonehenge	6	356
	74	Wood River	9	256
3AN	101	Terrell	1	339
	102	Lake Johnston	4	274
	103	Sutton	N/A	283
	134	Shamrock	4	358
	164	Chaplin	N/A	307
	193	Eyebrow	N/A	324
	194	Enfield	NIL	366
3BS	17	Val Marie	36	310
	45	Mankota	11	240
	75 A	Pinto Creek	20	271
	75 B	Pinto Creek	34	274
	76 A	Auvergne	22	227
	76 B	Auvergne	20	217
	77	Wise Creek	20	215
	78 A	Grassy Creek	40	329
	78 B	Grassy Creek	N/A	272
	78 C	Grassy Creek	N/A	305
	105	Glenbain	33	264
	106	Whiska Creek	14	195
	107	Lac Pelletier	14	259
	108	Bone Creek	17	318
3BN	138	Webb	2	318
	165	Morse	N/A	NIL
	166	Excelsior	NIL	437
	167	Sask. Landing	1	364
	168	Riverside	NIL	334
	226	Victory	N/A	386
	229	Lacadena	45	364
	255	Coteau	N/A	NIL
	257	Monet	50	394

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
4A	49	White Valley	N/A	300
	49 A	White Valley	6	238
	51	Reno	11.4	275
	79	Arlington	N/A	333
	109	Carmichael	24	404
	110	Popot	19	349
	111	Maple Creek	N/A	343
4B	139	Gull Lake	17	418
	141	Big Stick	11	389
	142	Enterprise	N/A	411
	169	Pitville	N/A	314
	171	Fox Valley	3	338
	229	Mry Creek	NIL	420
	231	Happyland	19	393
5A	181	Langenberg	N/A	322
	183	Fertile Belt	10	361
	185	Mcleod	9	336
	186	Abernethy	N/A	279
	211	Churchbridge	5	347
	213	Saltcoats	14	333
	216	Tullymet	N/A	219
	241	Calder	1	318
	243	Wallace	N/A	335
	245 A	Garry	13	407
	245 B	Garry	5	420
	246	Ituna Bon Accord	10	283
	247	Kelross	5	411
	248	Touchwood	NIL	241
5B	271	Cote	15	321
	275	Insinger	N/A	NIL
	276	Foam Lake	N/A	537
	277	Emerald	NIL	513
	278	Kutawa	N/A	NIL
	305	Invermay	3	660
	307	Elfros	12	613
	308	Big Quill	16	406
	333	Clayton	N/A	133
	335	Hazel Dell	N/A	377
	336	Sasman	28	494
	337	Lakeview	8	401
	338	Lakeside	6	472
	366	Kelvington	12	345
6A	190 A	Dufferin	N/A	243
	190 B	Dufferin	10	339
	190 C	Dufferin	NIL	268
	190 D	Dufferin	N/A	314
	219 A	Longlaketon	N/A	250
	219 B	Longlaketon	9	293
	220	Micklep	8	306
	221	Sarnia	N/A	369
	222	Craik	34	401
	251	Big Arm	6	372
	252	Arm River	2	323
	279	Mount Hope	NIL	354
	280	Wreford	N/A	323
	282	McCranney	8	400
	309	Prairie Rose	11	407
	313	Last River	N/A	306
	339	Lemy	6.6	460
	340	Wolverine	1	461
	341	Viscount	30	429
	343	Blucher	2.5	445
6B	253	Willner	N/A	310
	283	Rosedale	N/A	409
	284	Rudy	5	438
	285	Fertile Valley	11	372
	286	Milden	30	391
	314	Dundurn	10	444
	344	Corman Park	2	430
	345	Perdue	18	484
	376 A	Eagle Creek	N/A	550
	376 B	Eagle Creek	1	443
7A	260	Newcombe	N/A	346
	260 B	Newcombe	21	264
	287	St. Andrews	17	320
	288	Pleasant Valley	22	349
	290 A	Kindersley	2	261
	290 B	Kindersley	N/A	306
	292	Milton	N/A	280
	317 A	Marriott	21	451
	317 B	Marriott	17	449
	318 A	Mountain View	25	380
	318 B	Mountain View	47	479
	320	Oakdale	N/A	286
	321	Prairiedale	5	331

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
7B	347 A	Biggar	10	373
	347 B	Biggar	N/A	415
	350 A	Manpoca	29	365
	350 B	Manpoca	29	319
	351	Progress	33	368
	352	Heart's Hill	10	463
	377	Glenside	2	430
	378 A	Rosemount	15	336
	378 B	Rosemount	2	400
	379	Reford	22	371
	380	Tramping Lake	2	329
	381	Grass Lake	N/A	396
	382	Eye Hill	N/A	336
	409	Buffalo	2	455
	410	Round Valley	51	409
	411	Senlac	10	365
8A	396	Porcupine	N/A	346
	397	Barrier Valley	8.5	428
	428	Star City	N/A	336
	456	Arborfield	3	340
	457	Connaught	12	499
	458	Willow Creek	3	421
	488	Moose Range	8	368
8B	369	St. Peter	5	426
	370	Humboldt	30	527
	371	Bayne	6	452
	372	Grant	28.9	419
	402	Fish Creek	11	368
	429 A	Flett's Springs	N/A	254
	429 B	Flett's Springs	10	402
	430	Invergordon	N/A	43
	459	Kinistino	N/A	388
8AE	461	Prince Albert	0.3	404
	489	Torch River	3	409
	520	Paddockwood	N/A	67
8AW	406	Mayfield	NIL	314
	435	Radberry	N/A	406
	436	Douglas	N/A	241
	463	Duck Lake	20	334
	464	Leask	32	421
	467	Round Hill	N/A	NIL
	494	Canwood	N/A	413
	496	Spiritwood	13	440
	497	Medstead	NIL	291
9B	438	Battle River	N/A	358
	440	Hillsdale	N/A	325
	442	Manitou Lake	4.6	384
	498 A	Parkdale	NIL	325
	498 B	Parkdale	N/A	284
	499	Marvin	1	273
	501 A	Franchman Butte	4	288
	501 B	Franchman Butte	2	277
	501 C	Franchman Butte	4	322
	502	Britannia	12	340
	561	Loon Lake	N/A	290
	588 A	Meadow Lake	4	320
	588 B	Meadow Lake	13	345
	588 C	Meadow Lake	NIL	325
	588 D	Meadow Lake	5	330
	622	Beaver River	N/A	299

Municipality No: A, B, C, and D - more than one reporter

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Cumulative Rainfall

From: April 1, 2010
To: August 2, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 4, 2010

Cropland Topsoil Moisture Conditions

August 3, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 4, 2010

Cumulative Rainfall

From: April 1, 2010
To: August 2, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 4, 2010

Cropland Topsoil Moisture Conditions

August 3, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 4, 2010

Hay and Pasture Topsoil Moisture Conditions

August 3, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 4, 2010

Hay and Pasture Topsoil Moisture Conditions

August 3, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 4, 2010